


AMENDMENTS TO THE CLAIMS:

This listing of claims will replace all prior versions and listings of claims in the application:

1-56. (Canceled)

57. (Currently Amended) A cosmetic or pharmaceutical composition comprising, in a physiologically acceptable medium, at least one polymer comprising at least one monomeric compound of formula (I):




wherein:


- R₂ and R₃, which are present on the same ring or each on a different ring, are chosen from, independently of each other, hydrogen, halogens, and groups of formula -X-G-P (II), with the proviso that at least one of the radicals R₂ and R₃ is chosen from groups of formula (II), wherein:

- X is chosen from the groups -O-, -S-, -SO-, -SO₂-, -NH-, and -NR-, wherein R is chosen from linear, branched, and cyclic, saturated and unsaturated ~~carbon-based~~ radicals containing 1 to 30 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;


- G is chosen from linear, branched, and cyclic, saturated and unsaturated divalent ~~carbon-based~~ radicals containing 1 to 32 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;
- P is a polymerizable group chosen from formulae (IIIa) to (IIIc):



(IIIa)




(IIIb)



(IIIc)

wherein:

- R' is chosen from H, and linear and branched saturated C₁₋₆ ~~hydrocarbon-based~~ radicals;
- X' is chosen from O, NH, and NR'' with R'' being a radical chosen from C₁₋₆ alkyl, C₆₋₁₀ aryl, (C₆₋₁₀)aryl(C₁₋₆)alkyl, and (C₁₋₆)alkyl(C₆₋₁₀)aryl radicals, the alkyl and/or aryl groups optionally being substituted with at least one group chosen from OH, halogen, C₁₋₆ alkoxy and C₆₋₁₀ aryloxy;
- m is equal to 0 or 1; n is equal to 0 or 1; p is equal to 0, 1 or 2; and
- B is chosen from the divalent aromatic groups (IVa) to (IVd):



wherein:

- R1 is chosen from linear, branched, and cyclic, saturated and unsaturated ~~carbon-based~~ radicals containing 1 to 32 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;

- R22 is chosen from a hydrogen atom and linear, branched, and/ cyclic, saturated and unsaturated ~~carbon-based~~ radicals containing 1 to 32 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;

- R20 and R21 are, independently of each other, chosen from a hydrogen atom, linear and branched C₁₋₈ alkyl radicals, and cyclopentyl, cyclohexyl, cyclooctyl, cyclodecyl, cyclododecyl, benzyl, naphthyl and phenyl radicals.

58. (Previously Presented) The composition according to claim 57, wherein R₂ is hydrogen and R₃ is a group of formula (II).

59. (Currently Amended) The composition according to claim 57, wherein X is chosen from -O-, -NH- and -NR-, wherein R is chosen from linear, branched, and cyclic, saturated and unsaturated ~~hydrocarbon-based~~ radicals optionally


comprising a hydrocarbon-based ring chosen from saturated and unsaturated rings, containing 2 to 18 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂, and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S.

60. (Previously Presented) The composition according to claim 59, wherein X is chosen from -NH- and -NR-, and wherein R is a cyclohexyl radical.

61. (Currently Amended) The composition according to claim 57, wherein G is chosen from linear, branched, and cyclic, saturated and unsaturated divalent ~~hydrocarbon-based~~ radicals optionally comprising a hydrocarbon-based ring chosen from saturated and unsaturated rings, containing 2 to 18 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂, and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, and Si.

62. (Currently Amended) The composition according to claim 57, wherein G is chosen from linear and branched, saturated divalent ~~hydrocarbon-based~~ radicals optionally comprising a saturated hydrocarbon-based ring containing 2 to 18 carbon atoms.


63. (Previously Presented) The composition according to claim 57, wherein P is chosen from the following formulae:




and wherein R' is chosen from H and methyl.

64. (Currently Amended) The composition according to claim 57, wherein B is chosen from groups of formula (IVa) wherein R1 is chosen from linear, branched, and cyclic, saturated carbon-based radicals containing 1 to 32 carbon atoms.

65. (Previously Presented) The composition according to claim 57, wherein the monomeric compound is chosen from the following formulae wherein R is chosen from H and methyl:





66. (Previously Presented) The composition according to claim 57, wherein the at least one polymer is chosen from homopolymers of the at least one monomeric compound of formula (I).

67. (Previously Presented) The composition according to claim 57, wherein the at least one polymer is chosen from copolymers comprising only monomeric compounds of formula (I).


68. (Previously Presented) The composition according to claim 57, wherein the at least one polymer is chosen from copolymers comprising at least one monomeric compound of formula (I) and at least one additional comonomer.

69. (Previously Presented) The composition according to claim 67,


wherein the at least one polymer is chosen from statistical, alternating, grafted, block, and gradient copolymers.

70. (Previously Presented) The composition according to claim 68, wherein the at least one monomeric compound of formula (I) is present in an amount ranging from 0.01% to 70% by weight relative to the weight of the at least one polymer, and wherein the at least one additional comonomer comprises the remaining weight percent for a total polymer weight percent of 100%.

71. (Currently Amended) The composition according to claim 68, wherein the at least one additional comonomer is chosen from compounds of formula (A) and formula (B):



(A)



(B)

wherein:

- Ra₁ is chosen from linear, branched, and cyclic, saturated and unsaturated ~~carbon-based~~ radicals containing 1 to 32 carbon atoms; optionally substituted with at least one group chosen from =O, OH, NH₂, and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;

- Rb₁ is chosen from hydrogen, halogen atoms, linear, branched and cyclic, saturated and unsaturated ~~carbon-based~~ radicals containing 1 to 12 carbon atoms,


optionally substituted with at least one group chosen from =O, OH and NH₂ and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S; groups NRR' wherein R and R' are chosen, independently of each other, from hydrogen and linear, cyclic and branched, saturated C₁₋₆ ~~hydrocarbon-based~~ radicals;

- Ra₂ and Ra₃, which are present on the same ring or each on a different ring, are chosen, independently of each other, from hydrogen, halogens, and groups of formula -Xa-Ga-Pa (II), with the proviso that at least one of the radicals Ra₂ and Ra₃ is chosen from groups of formula (II), wherein:


- Xa is chosen from -O-, -S-, -SO-, -SO₂-, -NH-, and -NR₄- wherein R₄ is chosen from linear, branched and cyclic, saturated and unsaturated ~~carbon-based~~ radicals containing 1 to 30 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;

- Ga is chosen from linear, branched and cyclic, saturated and unsaturated divalent ~~carbon-based~~ radicals containing 1 to 32 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂, and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;


- Pa is a polymerizable group chosen from formulae (IIIa) to (IIIc):



(IIIa)



(IIIb)



(IIIc)

wherein:

- R' is chosen from H and linear and branched, saturated C₁₋₆ ~~hydrocarbon-based~~ radicals;


- X' is chosen from O, NH and NR'' with R'' being a radical chosen from C₁₋₆ alkyl, C₆₋₁₀ aryl, (C₆₋₁₀)aryl(C₁₋₆)alkyl, and (C₁₋₆)alkyl(C₆₋₁₀)aryl radicals, the alkyl and/or aryl groups optionally being substituted with at least one group chosen from OH, halogens, C₁₋₆ alkoxy, and C₆₋₁₀ aryloxy groups; and

- m is equal to 0 or 1; n is equal to 0 or 1; p is equal to 0, 1 or 2.

72. (Previously Presented) The composition according to claim 68, wherein the at least one additional comonomer is chosen from the monomers (i) to (viii):

(i) ethylenic hydrocarbons containing from 2 to 10 carbons;

(ii) (meth)acrylates chosen from:



wherein R'₃ is chosen from:

- linear and branched alkyl groups containing from 1 to 18 carbon atoms, optionally intercalated with at least one heteroatom chosen from O, N, S and P; said alkyl groups optionally being substituted with at least one substituent chosen from hydroxyl groups, halogen atoms, and groups Si(R₄R₅), wherein R₄ and R₅, which may be identical or different, are chosen from C₁₋₆ alkyl groups and phenyl groups;

- C₃ to C₁₂ cycloalkyl groups;


- C₃ to C₂₀ aryl groups;

- C₄ to C₃₀ aralkyl groups (C₁ to C₈ alkyl groups)
- 4- to 12-membered heterocyclic groups comprising at least one heteroatom chosen from O, N and S, the ring being chosen from aromatic and non-aromatic rings;
- heterocycloalkyl groups (1 to 4 C alkyls)

said cycloalkyl, aryl, aralkyl, heterocyclic and heterocycloalkyl groups being optionally substituted with at least one substituent chosen from hydroxyl groups, halogen atoms, and linear and branched C₁₋₄ alkyl groups, optionally intercalated with at least one heteroatom chosen from O, N, S and P, said alkyl groups optionally substituted with at least one substituent chosen from hydroxyl groups, halogen atoms, and groups Si(R₄R₅), wherein R₄ and R₅, which may be identical or different, are chosen from C₁ to C₆ alkyl groups and phenyl groups, and

- groups -(C₂H₄O)_m-R'', wherein m = 5 to 150 and R'' is chosen from H and C₁ to C₃₀ alkyl groups;

(iii) (meth)acrylamides of formula:



wherein:

- R₈ is chosen from H and methyl; and
- R₇ and R₆, which may be identical or different, are chosen from:
 - hydrogen;
 - linear and branched alkyl groups of 1 to 18 carbon atoms, optionally intercalated with at least one heteroatom chosen from O, N, S and P; said alkyl group

optionally being substituted with at least one substituent chosen from hydroxyl groups, halogen atoms, and groups $\text{Si}(\text{R}_4\text{R}_5)$, wherein R_4 and R_5 , which may be identical or different, are chosen from C_1 to C_6 alkyl groups and phenyl groups;

- C_3 to C_{12} cycloalkyl groups;
- C_3 to C_{20} aryl groups;
- C_4 to C_{30} aralkyl groups (C_1 to C_8 alkyl groups)
- 4- to 12-membered heterocyclic groups containing at least one heteroatom chosen from O, N and S, the ring being chosen from aromatic and non-aromatic; and
- heterocycloalkyl groups (1 to 4 C alkyls),

said cycloalkyl, aryl, aralkyl, heterocyclic and heterocycloalkyl groups being optionally substituted with at least one substituent chosen from hydroxyl groups, halogen atoms, and linear and branched C_1 - C_4 alkyl groups, optionally intercalated with at least one heteroatom chosen from O, N, S and P, said alkyl groups optionally being substituted with at least one substituent chosen from hydroxyl groups, halogen atoms and groups $\text{Si}(\text{R}_4\text{R}_5)$, wherein R_4 and R_5 , which may be identical or different, are chosen from C_1 to C_6 alkyl groups and phenyl groups;

(iv) vinyl compounds chosen from formulae:

$\text{CH}_2=\text{CH}-\text{R}_9$, $\text{CH}_2=\text{CH}-\text{CH}_2-\text{R}_9$ and $\text{CH}_2=\text{C}(\text{CH}_3)-\text{CH}_2-\text{R}_9$,

wherein:

- R_9 is chosen from hydroxyl groups, halogens, NH_2 , OR_{10} wherein R_{10} is chosen from phenyl groups, and C_1 to C_{12} alkyl groups; acetamide (NHCOCH_3); groups OCOR_{11} wherein R_{11} is chosen from linear and branched alkyl groups of 2 to 12 carbons; and groups chosen from:

- linear and branched alkyl groups of 1 to 18 carbon atoms, optionally intercalated with at least one heteroatom chosen from O, N, S and P, said alkyl group being optionally substituted with at least one substituent chosen from hydroxyl groups, halogen atoms, and groups $\text{Si}(\text{R}_4\text{R}_5)$, wherein R_4 and R_5 , which may be identical or different, are chosen from C_1 to C_6 alkyl groups and phenyl groups;

- C_3 to C_{12} cycloalkyl groups;

- C_3 to C_{20} aryl groups;

- C_4 to C_{30} aralkyl groups (C_1 to C_8 alkyl groups)

- 4- to 12-membered heterocyclic groups comprising at least one heteroatom chosen from O, N and S, the ring being chosen from aromatic and non-aromatic rings;

- heterocycloalkyl groups (1 to 4 C alkyls) optionally substituted with at least one substituent chosen from hydroxyl groups, halogen atoms, and linear and branched C_1 to C_4 alkyl groups, optionally intercalated with at least one heteroatom chosen from O, N, S and P, said alkyl groups being optionally substituted with at least one substituent chosen from hydroxyl groups, halogen atoms, and groups $\text{Si}(\text{R}_4\text{R}_5)$ wherein R_4 and R_5 , which may be identical or different, are chosen from C_1 to C_6 alkyl groups and phenyl groups;

(v) (meth)acrylic, (meth)acrylamide, and vinyl monomers comprising at least one group chosen from fluoro and perfluoro groups;

(vi) silicone-based (meth)acrylic, (meth)acrylamide, and vinyl monomers;

(vii) ethylenically unsaturated monomers comprising at least one functional group chosen from carboxylic acid, phosphoric acid, sulfonic acid, anhydride, and salts

thereof; and

(viii) ethylenically unsaturated monomers comprising at least one tertiary amine functional group and the salts thereof.

73. (Previously Presented) The composition according to claim 68, wherein the at least one additional comonomer is present in an amount ranging from 30% to 99.99% by weight relative to the weight of the at least one polymer.

74. (Previously Presented) The composition according to claim 68, wherein the at least one additional comonomer is chosen from C₁-C₁₈ alkyl and C₃-C₁₂ cycloalkyl (meth)acrylates.

75. (Previously Presented) The composition according to claim 74, wherein the at least one additional comonomer is chosen from methyl acrylate, methyl methacrylate, isobornyl acrylate, isobornyl methacrylate, isobutyl acrylate, isobutyl methacrylate, 2-ethylhexyl acrylate, 2-ethylhexyl methacrylate, dodecyl acrylate, dodecyl methacrylate, stearyl acrylate, stearyl methacrylate, trifluoroethyl acrylate, trifluoroethyl methacrylate, acrylic acid, methacrylic acid, methacryloxypropyltris(trimethylsiloxy)silane, acryloxypropyltris(trimethylsiloxy)silane, acryloxypropylpolydimethylsiloxane, and methacryloxypropylpolydimethylsiloxane.

76. (Previously Presented) The composition according to claim 57, wherein the polymer has a weight-average molecular mass (M_w) ranging from 5,000 to 600,000 g/mol.

77. (Previously Presented) The composition according to claim 57, wherein the at least one polymer is present, alone or as a mixture, in an amount ranging from 0.01% to 60% by weight relative to the total weight of the composition.

78. (Previously Presented) The composition according to claim 57, wherein the physiologically acceptable medium comprises at least one hydrophilic medium chosen from water and a mixture of water and at least one hydrophilic organic solvent; and/or comprises at least one fatty phase.

79. (Previously Presented) The composition according to claim 78, wherein the at least one fatty phase is chosen from waxes, pasty fatty substances, gums, lipophilic organic solvents, and oils.

80. (Previously Presented) The composition according to claim 57, further comprising at least one particulate phase chosen from pigments, nacles, and fillers.

81. (Previously Presented) The composition according to claim 57, further comprising at least one dyestuff chosen from water-soluble dyes and liposoluble dyes.

82. (Previously Presented) The composition according to claim 57, comprising at least one additional polymer.

83. (Previously Presented) The composition according to claim 82, wherein the at least one additional polymer is a film forming polymer.

84. (Previously Presented) The composition according to claim 57, further comprising at least one adjuvant chosen from vitamins, thickeners, gelling agents, trace elements, softeners, sequestrants, fragrances, acidifying agents, basifying agents, preserving agents, sunscreens, surfactants, antioxidants, hair-loss counteractants, antidandruff agents, propellants, and ceramides.


85. (Previously Presented) The composition according to claim 57, wherein the composition is in a form chosen from suspensions, dispersions, optionally thickened and gelled oily solutions, oil-in-water emulsions, water-in-oil emulsions,

multiple emulsions, gels, mousses, oily gels, emulsified gels, dispersions of vesicles, two-phase lotions, multi-phase lotions, sprays, loose powders, compact powders, cast powders, anhydrous pastes, lotions, creams, pomades, soft pastes, ointments, cast solids, molded solids, and compacted solids.

86. (Previously Presented) The composition according to claim 57, wherein the composition is in the form of care and makeup products for bodily and facial skin, the lips, the nails, the eyelashes, the eyebrows, and the hair; anti-sun and self-tanning products; and hair products for caring for, treating, shaping, making up, and dyeing the hair.

87. (Previously Presented) The composition according to claim 57, wherein the composition is in the form of makeup compositions, lip products, concealer products, blushers, mascara, eyeliners, eyebrow makeup products, lip pencils, eye pencils, nail products, nailcare products, body makeup products, hair makeup products, compositions for protecting and caring for the skin of the face, the neck, the hands, and the body, anti-wrinkle compositions, moisturizing and treating compositions, anti-sun compositions; artificial tanning compositions; and hair products.

88. (Withdrawn - Currently Amended) A cosmetic process for making up and/or caring for keratin materials comprising applying to said materials of at least one composition comprising, in a physiologically acceptable medium, at least one polymer comprising at least one monomeric compound of formula (I):




wherein:

- R_2 and R_3 , which are present on the same ring or each on a different ring, are chosen from, independently of each other, hydrogen, halogens, and groups of formula -X-G-P (II), with the proviso that at least one of the radicals R_2 and R_3 is chosen from groups of formula (II), wherein:


- X is chosen from the groups -O-, -S-, -SO-, -SO₂-, -NH-, and -NR-, wherein R is chosen from linear, branched, and cyclic, saturated and unsaturated ~~carbon-based~~ radicals containing 1 to 30 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;

- G is chosen from linear, branched, and cyclic, saturated and unsaturated divalent ~~carbon-based~~ radicals containing 1 to 32 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;


- P is a polymerizable group chosen from formulae (IIIa) to (IIIc):



(IIIa)



(IIIb)



(IIIc)


wherein:

- R' is chosen from H, and linear and branched saturated C₁₋₆ ~~hydrocarbon-~~
~~based-radicals~~;


- X' is chosen from O, NH, and NR'' with R'' being a radical chosen from C₁₋₆
alkyl, C₆₋₁₀ aryl, (C₆₋₁₀)aryl(C₁₋₆)alkyl, and (C₁₋₆)alkyl(C₆₋₁₀)aryl radicals, the alkyl and/or
aryl groups optionally being substituted with at least one group chosen from OH,
halogen, C₁₋₆ alkoxy and C₆₋₁₀ aryloxy;

- m is equal to 0 or 1; n is equal to 0 or 1; p is equal to 0, 1 or 2; and


- B is chosen from the divalent aromatic groups (IVa) to (IVd):




(IVa)



(IVb)



(IVc)



(IVd)

wherein:


- R1 is chosen from linear, branched, and cyclic, saturated and unsaturated
~~carbon-based-radicals~~ containing 1 to 32 carbon atoms, optionally substituted with at

least one group chosen from =O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;

- R22 is chosen from a hydrogen atom and linear, branched, and/ cyclic, saturated and unsaturated ~~carbon-based~~ radicals containing 1 to 32 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;

- R20 and R21 are, independently of each other, chosen from a hydrogen atom, linear and branched C₁₋₈ alkyl radicals, and cyclopentyl, cyclohexyl, cyclooctyl, cyclodecyl, cyclododecyl, benzyl, naphthyl and phenyl radicals.

89. (Withdrawn - Currently Amended) A monomeric compound of formula (I):



wherein:


- R₂ and R₃, which are present on the same ring or each on a different ring, are chosen from, independently of each other, hydrogens, halogens, and groups of formula -X-G-P (II), with the proviso that at least one of the radicals R₂ and R₃ is chosen from groups of formula (II), wherein:

- X is chosen from the groups -O-, -S-, -SO-, -SO₂-, -NH-, and -NR-, wherein R is chosen from linear, branched and cyclic, saturated and unsaturated ~~carbon-based~~


radicals containing 1 to 30 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;

- G is chosen from linear, branched and cyclic, saturated and unsaturated divalent ~~carbon-based~~ radicals containing 1 to 32 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;


- P is a polymerizable group chosen from the formulae (IIIa) to (IIIc):



(IIIa)



(IIIb)



(IIIc)


wherein:

- R' is chosen from H, and linear and branched, saturated C₁₋₆ ~~hydrocarbon-based~~ radicals;

- X' is chosen from O, NH, and NR'' with R'' being a radical chosen from C₁₋₆ alkyl, C₆₋₁₀ aryl, (C₆₋₁₀)aryl(C₁₋₆)alkyl, and (C₁₋₆)alkyl(C₆₋₁₀)aryl radicals, the alkyl and/or aryl groups being optionally substituted with at least one group chosen from OH, halogen, C₁₋₆ alkoxy, and C₆₋₁₀ aryloxy;

- m is equal to 0 or 1; n is equal to 0 or 1; p is equal to 0, 1 or 2; and

- B is chosen from the divalent aromatic groups (IVa) to (IVd):



wherein:

- R1 is chosen from linear, branched and cyclic, saturated and unsaturated ~~carbon-based~~ radicals containing 1 to 32 carbon atoms, optionally substituted with at least one group chosen from = O, OH, NH₂ and halogen atoms;

- R22 is chosen from a hydrogen atom and linear, branched and cyclic, saturated and unsaturated ~~carbon-based~~ radicals containing 1 to 32 carbon atoms, optionally substituted with at least one group chosen from = O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;

- R20 and R21 are chosen from, independently of each other, a hydrogen atom, linear and branched C₁₋₈ alkyl radicals and cyclopentyl, cyclohexyl, cyclooctyl, cyclodecyl, cyclododecyl, benzyl, naphthyl, and phenyl radicals; with the exclusion of compounds for which, simultaneously, P is of formula (IIIa), X' is O, m = 1, X is NH, and B is of formula (IVc).

90. (Withdrawn) The monomeric compound according to claim 89, wherein R₂ is a hydrogen atom and R₃ is a group of formula (II).

91. (Withdrawn - Currently Amended) The monomeric compound according to claim 89, wherein X is chosen from -O-, -NH-, and -NR- with R being chosen from

linear, branched and cyclic, saturated and unsaturated ~~hydrocarbon-based-radicals~~ optionally comprising a hydrocarbon-based ring that is itself chosen from saturated and unsaturated rings, containing 2 to 18 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂, and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si, and S.

92. (Withdrawn) The monomeric compound according to claim 91, wherein R is chosen from ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, pentyl, hexyl, cyclohexyl, octyl, cyclooctyl, decyl, cyclodecyl, dodecyl, cyclododecyl, phenyl, and benzyl radicals.

93. (Withdrawn) The monomeric compound according to claim 91, wherein X is chosen from -NH- and -NR- wherein R is a cyclohexyl.


94. (Withdrawn - Currently Amended) The monomeric compound according to claim 89, wherein G is chosen from linear, branched and cyclic, saturated and unsaturated divalent ~~hydrocarbon-based-radicals~~ optionally comprising a hydrocarbon-based ring that is itself chosen from saturated and unsaturated rings, containing in total 2 to 18 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, and Si.

95. (Withdrawn - Currently Amended) The monomeric compound according to claim 94, wherein G is chosen from linear and branched, saturated divalent ~~hydrocarbon-based-radicals~~ optionally comprising a saturated hydrocarbon-based ring, containing in total 2 to 18 carbon atoms.

96. (Withdrawn) The monomeric compound according to claim 94, wherein

G is chosen from ethylene, n-propylene, isopropylene, 1-methylethylene, 2-methylethylene, n-butylene, isobutylene, pentylene, hexylene, cyclohexylene, heptylene, octylene, cyclooctylene, decylene, cyclodecylene, cyclohexyldimethylene, dodecylene, and cyclododecylene radicals.


97. (Withdrawn) The monomeric compound according to claim 89, wherein the polymerizable group P is chosen from the formulae:




wherein R' is chosen from H and methyl.


98. (Withdrawn - Currently Amended) The monomeric compound according to claim 89, wherein B is chosen from groups of formula (IVa) wherein R_1 is chosen from linear, branched and cyclic, saturated ~~carbon-based~~ radicals containing 1 to 32 carbon atoms.

99. (Withdrawn) The monomeric compound according to claim 89, chosen from the following formulae, wherein R is chosen from H and methyl:





100. (Currently Amended) A polymer comprising at least one monomeric compound of formula (I):



wherein:


- R_2 and R_3 , which are present on the same ring or each on a different ring, are chosen from, independently of each other, hydrogen, halogens, and groups of formula

-X-G-P (II), with the proviso that at least one of the radicals R_2 and R_3 is chosen from groups of formula (II), wherein:


- X is chosen from the groups -O-, -S-, -SO-, -SO₂-, -NH-, and -NR-, wherein R is chosen from linear, branched, and cyclic, saturated and unsaturated ~~carbon-based~~ radicals containing 1 to 30 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;

- G is chosen from linear, branched, and cyclic, saturated and unsaturated divalent ~~carbon-based~~ radicals containing 1 to 32 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;


- P is a polymerizable group chosen from formulae (IIIa) to (IIIc):



(IIIa)



(IIIb)




(IIIc)

wherein:

- R' is chosen from H, and linear and branched saturated C₁₋₆ ~~hydrocarbon-based~~ radicals;

- X' is chosen from O, NH, and NR'' with R'' being a radical chosen from C₁₋₆ alkyl, C₆₋₁₀ aryl, (C₆₋₁₀)aryl(C₁₋₆)alkyl, and (C₁₋₆)alkyl(C₆₋₁₀)aryl radicals, the alkyl and/or aryl groups optionally being substituted with at least one group chosen from OH, halogen, C₁₋₆ alkoxy and C₆₋₁₀ aryloxy;

- m is equal to 0 or 1; n is equal to 0 or 1; p is equal to 0, 1 or 2; and
- B is chosen from the divalent aromatic groups (IVa) to (IVd):



wherein:

- R¹ is chosen from linear, branched, and cyclic, saturated and unsaturated ~~carbon-based~~ radicals containing 1 to 32 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;

- R²² is chosen from a hydrogen atom and linear, branched, and/ cyclic, saturated and unsaturated ~~carbon-based~~ radicals containing 1 to 32 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;

- R₂₀ and R₂₁ are, independently of each other, chosen from a hydrogen atom, linear and branched C₁₋₈ alkyl radicals, and cyclopentyl, cyclohexyl, cyclooctyl, cyclodecyl, cyclododecyl, benzyl, naphthyl and phenyl radicals.

101. (Previously Presented) The polymer according to claim 100, wherein said polymer is a homopolymer of the at least one monomeric compound.

102. (Previously Presented) The polymer according to claim 100, wherein

said polymer is a copolymer comprising only the at least one monomeric compound.


103. (Previously Presented) The polymer according to claim 102, wherein the at least one monomeric compound is present in an amount, for each monomeric compound, ranging from 0.5% to 99.5% by weight relative to the total weight of the polymer.

104. (Previously Presented) The polymer according to claim 100, wherein said polymer comprises a copolymer comprising the at least one monomeric compound and the at least one additional comonomer.


105. (Previously Presented) The polymer according to claim 100, wherein said polymer is a copolymer chosen from statistical, alternating, grafted, block and gradient copolymers.

106. (Previously Presented) The polymer according to claim 104, wherein the at least one monomeric compound is present in an amount ranging from 0.01% to 70% by weight relative to the weight of said polymer, and the at least one additional comonomer, alone or as a mixture, comprises the remaining weight percent of said polymer to 100% by weight.

107. (Currently Amended) The polymer according to claim 104, wherein the at least one additional comonomer is chosen from the compounds of formula (A) and of formula (B):



(A)



(B)

wherein:

- Ra_1 is chosen from linear, branched, and cyclic, saturated and unsaturated ~~carbon-based~~ radicals containing 1 to 32 carbon atoms; optionally substituted with at least one group chosen from =O, OH, NH_2 , and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;


- Rb_1 is chosen from hydrogen, halogen atoms, linear, branched and cyclic, saturated and unsaturated ~~carbon-based~~ radicals containing 1 to 12 carbon atoms, optionally substituted with at least one group chosen from =O, OH and NH_2 and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S; groups NRR' wherein R and R' are chosen, independently of each other, from hydrogen and linear, cyclic and branched, saturated C_{1-6} ~~hydrocarbon-based~~ radicals;

- Ra_2 and Ra_3 , which are present on the same ring or each on a different ring, are chosen, independently of each other, from hydrogen, halogens, and groups of formula -Xa-Ga-Pa (II), with the proviso that at least one of the radicals Ra_2 and Ra_3 is chosen from groups of formula (II), wherein:


- Xa is chosen from -O-, -S-, -SO-, -SO₂-, -NH-, and -NR₄- wherein R₄ is chosen from linear, branched and cyclic, saturated and unsaturated ~~carbon-based~~ radicals containing 1 to 30 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;

- Ga is chosen from linear, branched and cyclic, saturated and unsaturated divalent ~~carbon-based~~ radicals containing 1 to 32 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂, and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;


- Pa is a polymerizable group chosen from formulae (IIIa) to (IIIc):



(IIIa)



(IIIb)



(IIIc)

wherein:

- R' is chosen from H and linear and branched, saturated C₁₋₆ ~~hydrocarbon-based~~ radicals;

- X' is chosen from O, NH and NR'' with R'' being a radical chosen from C₁₋₆ alkyl, C₆₋₁₀ aryl, (C₆₋₁₀)aryl(C₁₋₆)alkyl, and (C₁₋₆)alkyl(C₆₋₁₀)aryl radicals, the alkyl and/or aryl groups optionally being substituted with at least one group chosen from OH, halogens, C₁₋₆ alkoxy, and C₆₋₁₀ aryloxy groups; and

- m is equal to 0 or 1; n is equal to 0 or 1; p is equal to 0, 1 or 2.

108. (Previously Presented) The polymer according to claim 104, wherein the at least one additional comonomer is chosen from hydrophilic comonomers, and is present in an amount ranging from 1% to 99.99% by weight relative to the total weight of the copolymer.

109. (Previously Presented) The polymer according to claim 104, wherein the at least one additional comonomer is chosen from hydrophobic comonomers and is present in an amount ranging from 1% to 99.99% by weight relative to the total weight of the copolymer.

110. (Previously Presented) The polymer according to claim 104, comprising at least one comonomer chosen from the following monomers:

- (i) ethylenic hydrocarbons containing from 2 to 10 carbons; and
- (ii) (meth)acrylates chosen from formulae:



wherein R'₃ is chosen from:

- linear and branched alkyl groups containing from 1 to 18 carbon atoms, optionally intercalated with at least one heteroatom chosen from O, N, S and P; said alkyl groups optionally being substituted with at least one substituent chosen from hydroxyl groups, halogen atoms, and groups Si(R₄R₅), wherein R₄ and R₅, which may be identical or different, are chosen from C₁₋₆ alkyl groups and phenyl groups;

- C₃ to C₁₂ cycloalkyl groups;
- C₃ to C₂₀ aryl groups;

- C₄ to C₃₀ aralkyl groups (C₁ to C₈ alkyl groups)


- 4- to 12-membered heterocyclic groups comprising at least one heteroatom chosen from O, N and S, the ring being chosen from aromatic and non-aromatic rings;

- heterocycloalkyl groups (1 to 4 C alkyls)

said cycloalkyl, aryl, aralkyl, heterocyclic and heterocycloalkyl groups being optionally substituted with at least one substituent chosen from hydroxyl groups, halogen atoms, and linear and branched C₁₋₄ alkyl groups, optionally intercalated with at least one heteroatom chosen from O, N, S and P, said alkyl groups optionally substituted with at least one substituent chosen from hydroxyl groups, halogen atoms, and groups Si(R₄R₅), wherein R₄ and R₅, which may be identical or different, are chosen from C₁ to C₆ alkyl groups and phenyl groups, and

- groups -(C₂H₄O)_m-R'', wherein m = 5 to 150 and R'' is chosen from H and C₁ to C₃₀ alkyl groups;

(iii) (meth)acrylamides of formula:



wherein:

- R₈ is chosen from H and methyl; and

- R₇ and R₆, which may be identical or different, are chosen from:

- hydrogen;

- linear and branched alkyl groups of 1 to 18 carbon atoms, optionally intercalated with at least one heteroatom chosen from O, N, S and P; said alkyl group

optionally being substituted with at least one substituent chosen from hydroxyl groups, halogen atoms, and groups $\text{Si}(\text{R}_4\text{R}_5)$, wherein R_4 and R_5 , which may be identical or different, are chosen from C_1 to C_6 alkyl groups and phenyl groups;

- C_3 to C_{12} cycloalkyl groups;
- C_3 to C_{20} aryl groups;
- C_4 to C_{30} aralkyl groups (C_1 to C_8 alkyl groups)

- 4- to 12-membered heterocyclic groups containing at least one heteroatom chosen from O, N and S, the ring being chosen from aromatic and non-aromatic; and

- heterocycloalkyl groups (1 to 4 C alkyls),

said cycloalkyl, aryl, aralkyl, heterocyclic and heterocycloalkyl groups being optionally substituted with at least one substituent chosen from hydroxyl groups, halogen atoms, and linear and branched C_1 - C_4 alkyl groups, optionally intercalated with at least one heteroatom chosen from O, N, S and P, said alkyl groups optionally being substituted with at least one substituent chosen from hydroxyl groups, halogen atoms and groups $\text{Si}(\text{R}_4\text{R}_5)$, wherein R_4 and R_5 , which may be identical or different, are chosen from C_1 to C_6 alkyl groups and phenyl groups;

(iv) vinyl compounds chosen from formulae:

$\text{CH}_2=\text{CH}-\text{R}_9$, $\text{CH}_2=\text{CH}-\text{CH}_2-\text{R}_9$ and $\text{CH}_2=\text{C}(\text{CH}_3)-\text{CH}_2-\text{R}_9$,

wherein:

- R_9 is chosen from hydroxyl groups, halogens, NH_2 , OR_{10} wherein R_{10} is chosen from phenyl groups, and C_1 to C_{12} alkyl groups; acetamide (NHCOCH_3); groups OCOR_{11} wherein R_{11} is chosen from linear and branched alkyl groups of 2 to 12 carbons; and groups chosen from:

- linear and branched alkyl groups of 1 to 18 carbon atoms, optionally intercalated with at least one heteroatom chosen from O, N, S and P, said alkyl group being optionally substituted with at least one substituent chosen from hydroxyl groups, halogen atoms, and groups $\text{Si}(\text{R}_4\text{R}_5)$, wherein R_4 and R_5 , which may be identical or different, are chosen from C_1 to C_6 alkyl groups and phenyl groups;

- C_3 to C_{12} cycloalkyl groups;

- C_3 to C_{20} aryl groups;

- C_4 to C_{30} aralkyl groups (C_1 to C_8 alkyl groups)

- 4- to 12-membered heterocyclic groups comprising at least one heteroatom chosen from O, N and S, the ring being chosen from aromatic and non-aromatic rings;

- heterocycloalkyl groups (1 to 4 C alkyls) optionally substituted with at least one substituent chosen from hydroxyl groups, halogen atoms, and linear and branched C_1 to C_4 alkyl groups, optionally intercalated with at least one heteroatom chosen from O, N, S and P, said alkyl groups being optionally substituted with at least one substituent chosen from hydroxyl groups, halogen atoms, and groups $\text{Si}(\text{R}_4\text{R}_5)$ wherein R_4 and R_5 , which may be identical or different, are chosen from C_1 to C_6 alkyl groups and phenyl groups;

(v) (meth)acrylic, (meth)acrylamide, and vinyl monomers comprising at least one group chosen from fluoro and perfluoro groups;

(vi) silicone-based (meth)acrylic, (meth)acrylamide, and vinyl monomers;

(vii) ethylenically unsaturated monomers comprising at least one functional group chosen from carboxylic acid, phosphoric acid, sulfonic acid, anhydride, and salts

thereof ; and

(viii) ethylenically unsaturated monomers comprising at least one tertiary amine functional group and the salts thereof.


111. (Previously Presented) The polymer according to claim 104, wherein the at least one additional comonomer is present in an amount ranging from 30% to 99.99% by weight relative to the weight of said polymer.

112. (Previously Presented) The polymer according to claim 104, wherein the at least one additional comonomer is chosen from C₁-C₁₈ alkyl and C₃-C₁₂ cycloalkyl (meth)acrylates

113. (Previously Presented) The polymer according to claim 112, wherein the at least one additional comonomer is chosen from methyl acrylate, methyl methacrylate, isobornyl acrylate, isobornyl methacrylate, isobutyl acrylate, isobutyl methacrylate, 2-ethylhexyl acrylate, 2-ethylhexyl methacrylate, dodecyl acrylate, dodecyl methacrylate, stearyl acrylate, stearyl methacrylate, trifluoroethyl acrylate and trifluoroethyl methacrylate; or alternatively acrylic acid, methacrylic acid, methacryloxypropyltris(trimethylsiloxy)silane, acryloxypropyltris(trimethylsiloxy)silane, acryloxypropylpolydimethylsiloxane, and methacryloxypropylpolydimethylsiloxane.

114. (Previously Presented) The polymer according to claim 104, having a weight-average molecular mass (M_w) ranging from 5,000 to 600,000 g/mol.

115. (Withdrawn - Currently Amended) A method for giving optical effects to cosmetic and/or pharmaceutical compositions, comprising,
combining a physiologically acceptable medium, and
at least one polymer comprising at least one monomeric compound of formula (I):




wherein:

- R_2 and R_3 , which are present on the same ring or each on a different ring, are chosen from, independently of each other, hydrogen, halogens, and groups of formula -X-G-P (II), with the proviso that at least one of the radicals R_2 and R_3 is chosen from groups of formula (II), wherein:


- X is chosen from the groups -O-, -S-, -SO-, -SO₂-, -NH-, and -NR-, wherein R is chosen from linear, branched, and cyclic, saturated and unsaturated ~~carbon-based~~ radicals containing 1 to 30 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;

- G is chosen from linear, branched, and cyclic, saturated and unsaturated divalent ~~carbon-based~~ radicals containing 1 to 32 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;


- P is a polymerizable group chosen from formulae (IIIa) to (IIIc):



(IIIa)



(IIIb)



(IIIc)


wherein:

- R' is chosen from H, and linear and branched saturated C₁₋₆ ~~hydrocarbon-~~
~~based~~ radicals;


- X' is chosen from O, NH, and NR'' with R'' being a radical chosen from C₁₋₆ alkyl, C₆₋₁₀ aryl, (C₆₋₁₀)aryl(C₁₋₆)alkyl, and (C₁₋₆)alkyl(C₆₋₁₀)aryl radicals, the alkyl and/or aryl groups optionally being substituted with at least one group chosen from OH, halogen, C₁₋₆ alkoxy and C₆₋₁₀ aryloxy;

- m is equal to 0 or 1; n is equal to 0 or 1; p is equal to 0, 1 or 2; and


- B is chosen from the divalent aromatic groups (IVa) to (IVd):




(IVa)



(IVb)



(IVc)



(IVd)

wherein:

- R1 is chosen from linear, branched, and cyclic, saturated and unsaturated ~~carbon-based~~ radicals containing 1 to 32 carbon atoms, optionally substituted with at

least one group chosen from =O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;

- R22 is chosen from a hydrogen atom and linear, branched, and/ cyclic, saturated and unsaturated ~~carbon-based~~ radicals containing 1 to 32 carbon atoms, optionally substituted with at least one group chosen from =O, OH, NH₂ and halogen atoms; and/or optionally interrupted with at least one heteroatom chosen from O, N, P, Si and S;

- R20 and R21 are, independently of each other, chosen from a hydrogen atom, linear and branched C₁₋₈ alkyl radicals, and cyclopentyl, cyclohexyl, cyclooctyl, cyclodecyl, cyclododecyl, benzyl, naphthyl and phenyl radicals.